


# Research Notes

Arm & Hammer Animal and Food Production




## Finalyse decreased *E. coli* on live cattle hides in 45 minutes.

### STUDY OVERVIEW<sup>1</sup>

Using the spray system found in commercial beef-processing facilities, a study was performed to determine the efficacy of Finalyse™ on the hides of live cattle.

- 50 heifers weighing approximately 1,000 lbs. were used in the study
  - 10/day, 5/treatment group
- The left and right side of the rump (1,000 cm<sup>2</sup>) of live cattle were inoculated with *E. coli* O157:H7
- Study conducted over a 5-day period
  - Day 1 was excluded due to the low numbers of bacteria recovered on the control animals
- Samples taken 45 minutes post-inoculation to establish pre-treatment levels of *E. coli* O157:H7
- Cattle were treated using the Finalyse Application System™
  - Control water wash
  - Finalyse solution
- Samples were taken 45 minutes post-treatment



### RESULTS

- Cattle treated with Finalyse exhibited a statistically significant decrease ( $P < 0.01$ ) in *E. coli* O157:H7 compared to controls at 45 minutes post-treatment
- Finalyse caused, on average, a 0.85<sub>Log</sub> CFU/mL mean reduction in *E. coli* O157:H7 when directly comparing post-treatment values

Statistical analysis using pre-treatment Log values as a covariate found the post-treatment least square mean value decreased 1.03<sub>Logs</sub> compared to the control

**TABLE 1** Statistical results

	Post-treatment least square means, adjusted for pre-treatment values		P-value	
	Control	Finalyse	Baseline	Treatment
<b>Pooled days 2-5</b>	3.99	2.96	<0.01	<0.01



#ScienceHearted

AHfoodchain.com

<sup>1</sup> Data on file, 2011.