Research Notes P-91

Arm & Hammer Animal Nutrition

AVIATOR supplementation reduced degree of infection in commercial layer pullets artificially infected with *Salmonella enteritidis*.

OBJECTIVE

Determine if feeding AVIATOR[™] to commercial layer pullets either from 1 day to 17 weeks or from 10 weeks to 17 weeks would affect *Salmonella enteritidis* colonization of both the digestive tract (cecum) and ovaries.

MATERIALS AND METHODS

Two hundred and fifty (250) one-day old Hyline W-36 pullets were purchased from Hyline, Mansfield, Georgia. Each bird was in an individual cage, with forty-eight (48) pullets per treatment group. The following treatments were tested:


- T1: No additive, challenge
- T2: AVIATOR SCP, 100g/MT supplemented from day 1 to study termination, challenge
- T3: AVIATOR SCP, 100g/MT supplemented from week 10 to study termination, challenge
- T4: No additive, no challenge

All birds were fed a non-medicated commercial layer diet in mash form. At 16 weeks-of-age each bird was orally challenged with a nalidixic acid resistant strain of *Salmonella enteritidis* (1.8×10^9 cfu/ml). One week (seven days) post-challenge all birds were necropsied and ceca and ovary samples taken. *Salmonella* was isolated using selective media culturing, and identified and serogrouped using poly-O *Salmonella* specific antiserum. *Salmonella* were enumerated using the most probable number (MPN) method. Data were analyzed statistically, and *P*<0.05 was considered significant.

RESULTS

Ceca Salmonella Prevalence and MPN.

Salmonella was identified in all of the ceca samples from all three of the challenged treatment groups. The prevalence of ceca samples with Salmonella loads that exceeded one million cfu/g (the upper quantitative limits of the MPN assay) are summarized in Figure 1. AVIATOR supplementation from day one of age trended to reduce the number of hens with high (over one million cfu/g) cecal Salmonella enteritidis compared to untreated or AVIATOR supplementation from 10 weeks of age (*P*=0.06).




Quantitative results for the ceca Salmonella MPNs are summarized in Figure 2, and the distribution of MPNs is illustrated in Figure 3. Observations which exceeded the MPN threshold were censored in the quantitative Salmonella analysis. Both AVIATORTM supplementation treatments reduced Salmonella enteritidis by over one log compared to untreated control (P=0.056).


Ovary Salmonella Prevalence. There was no significant difference between challenged treatment groups with respect to ovary Salmonella prevalence (*P*=0.32). All except two of the layers had Salmonella MPNs that were below the quantitative detection limit of the MPN assay. Decreased Salmonella counts in the ovaries is not surprising because ovaries naturally tend to clear Salmonella infection rapidly.

CONCLUSIONS

- AVIATOR supplementation from day 1 of age trended to reduce the number of hens with high (over one million cfu/g) cecal *Salmonella enteritidis* in challenged layers compared to untreated or AVIATOR supplementation from 10 weeks of age (*P*=0.06).
- Both AVIATOR supplementation treatments reduced *Salmonella enteritidis* by over one log in challenge trials compared to untreated control (*P*=0.056).
- A higher proportion of hens receiving AVIATOR from day 1 were noted to have lower *S. enteritidis* counts in the ceca compared to hens fed control or AVIATOR from week 10.
- No difference between treatments on ovarian *Salmonella* colonization was noted.

FIGURE 2 AVIATOR reduced Salmonella log₁₀ MPN per gram in ceca samples by 1.2 log





Samples that exceeded the upper limits of the MPN assay were arbitrarily assigned an MPN value of approximately 1×10^6 MPN/g for graphical illustration.





Based on study conducted by Dr. Hofacre at Southern Poultry Research Inc., Athens, GA © 2017 Church & Dwight Co., Inc. ARM & HAMMER™, the ARM & HAMMER logo and AVIATOR™ are trademarks of Church & Dwight Co., Inc. AV2791-0517