

Swine: minimum recommended feeding rates.¹

Making the uncertain certain



Animal Nutrition



A-MAX™ is an all-natural yeast culture product that helps animals digest feed more efficiently, resulting in improved average daily gain.



BG-MAX™ delivers specially processed yeast culture with aluminosilicate to bind toxins in poor-quality feeds. It is an all-natural fermentation product that can be used at all stages of growth.



CELMANAX™ combines yeast culture with a unique blend of Refined Functional Carbohydrates™ (RFC™) that enhances the gut environment, prepares the immune system, binds mycotoxins and improves nutrient digestibility.



KULACTIC™ delivers whey metabolites and organic acids to boost energy and help lower stomach pH for improved performance and feed conversion.

	Swine (kg/MT)			Swine (lbs/ton)		
	Sow	Nursery	Grow/finish	Sow	Nursery	Grow/finish
A-MAX Standard	10	20	10	20	40	20
A-MAX Concentrate	5	10	5	10	20	10
A-MAX Xtra	2.5	5	2.5	5	10	5
A-MAX Ultra™	1.25	2.5	1.25	2.5	5	2.5
A-MAX SCP ²	0.2	0.2	0.2	0.4	0.4	0.4
BG-MAX	2	2	1	4	4	2
CELMANAX	2	2	2	4	4	4
CELMANAX SCP ²	0.2	0.2	0.2	0.4	0.4	0.4
KULACTIC	25	25	25	50	50	50
	ml/hd/day			ml/hd/day		
CELMANAX Liquid ²	6	2	6	6	2	6

Additional notes:

1. All feeding rates are for normal production situations. When additional stressors are present (heat stress, mycotoxin loads, pathogen challenges, etc.) doses may be increased on the recommendation of the nutritionist or veterinarian. Contact your nutritionist for your optimum feeding rates.
2. When included in TMR mixer SCP and Liquid products should be added as a premix or diluted to assure uniform distribution. For conversion purposes 1 ml of CELMANAX Liquid equals 1 gram.

To learn more, contact your nutritionist, veterinarian or Arm & Hammer Animal Nutrition representative or visit AHanimalnutrition.com.

Poultry: minimum recommended feeding rates.¹

Making the uncertain certain



Animal Nutrition

	CHICKENS						TURKEYS					
	Chickens (kg/MT)			Chickens (lbs/ton)			Turkeys (kg/MT)			Turkeys (lbs/ton)		
	Layer	Broiler	Broiler/ breeder	Layer	Broiler	Broiler/ breeder	Breeder	Poults	Grow/ finish	Breeder	Poults	Grow/ finish
A-MAX Standard	6	6	6	12	12	12	6	6	6	12	12	12
A-MAX Concentrate	3	3	3	6	6	6	3	3	3	6	6	6
A-MAX Xtra	1.5	1.5	1.5	3	3	3	1.5	1.5	1.5	3	3	3
A-MAX Ultra™	0.75	0.75	0.75	1.5	1.5	1.5	0.75	0.75	0.75	1.5	1.5	1.5
A-MAX SCP ²	0.05	0.05	0.05	0.1	0.1	0.1	0.05	0.05	0.05	0.1	0.1	0.1
AVIATOR	0.5	0.5	0.5	1	1	1	0.5	0.5	0.5	1	1	1
AVIATOR SCP ²	0.05	0.05	0.05	0.1	0.1	0.1	0.05	0.05	0.05	0.1	0.1	0.1
BG-MAX	1	1	1	2	2	2	1	1	1	2	2	2
CELMANAX	0.5	0.5	0.5	1	1	1	0.5	0.5	0.5	1	1	1
CELMANAX SCP ²	0.05	0.05	0.05	0.1	0.1	0.1	0.05	0.05	0.05	0.1	0.1	0.1
KULACTIC	15	15	15	30	30	30	15	15	15	30	30	30
	ml/L						ml/L					
AVIATOR Liquid ²	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25
CELMANAX Liquid ²	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25



Additional notes:

1. All feeding rates are for normal production situations. When additional stressors are present (heat stress, mycotoxin loads, pathogen challenges, etc.) doses may be increased on the recommendation of the nutritionist or veterinarian. Contact your nutritionist for your optimum feeding rates.
2. When included in TMR mixer, SCP and Liquid products should be added as a premix or diluted to assure uniform distribution. For conversion purposes 1 ml of CELMANAX Liquid or AVIATOR Liquid equals 1 gram.

To learn more, contact your nutritionist, veterinarian or Arm & Hammer Animal Nutrition representative or visit AHanimalnutrition.com.